


Algorithm for High-Risk Well-Appearing Febrile Infant: 61-90 days of age, temp $\geq 38\text{C}^*$ **Footnotes:**

* The AAP does not include infants age > 60 days
† If observed at home, then recommend oral antimicrobials to treat presumed UTI if UA positive. Clinicians may manage infants at home if ALL following criteria are met: **1) Verbal teaching and written instructions** have been provided for monitoring throughout the period at home **2) Follow-up plans for re-evaluation** have been developed and are in place **3) Plans have been developed and are in place in case of change in clinical status**, including means of communication between family and providers and access to emergency medical care.

Algorithm for High-Risk Well-Appearing Febrile Infant: 61-90 days of age, temp \geq 38C*

Exclusion criteria

1. Infants who have received immunizations within the last 48 hrs
2. Infants with clinical bronchiolitis, with or without positive tests for RSV
3. Infants with COVID-19



Consult Pediatrics

Abbreviations

1. Cath: catheter
2. CBC: complete blood count
3. LP: lumbar puncture
4. RSV: respiratory syncytial virus
5. UA: urinalysis
6. UCx: urine culture
7. UTI: urinary tract infection
8. Pabx: parenteral antibiotics

Inclusion criteria

Infants 61-90 days of age*, well-appearing, temp \geq 38C AND one of the following:

1. Premature infants < 37 wks gestational age
2. Infants with documented or suspected immune compromise
3. Infants whose neonatal course was complicated by surgery or infection.
4. Infants with congenital or chromosomal abnormalities
5. Medically fragile infants requiring some form of technology or ongoing therapeutic intervention to sustain life

CA FIRST Protocol v2.1

Supplemental Information

Footnotes

* The AAP does not include infants age > 60 days
† If observed at home, then recommend oral antimicrobials to treat presumed UTI if UA positive. Clinicians may manage infants at home if ALL following criteria are met: **1) Verbal teaching and written instructions** have been provided for monitoring throughout the period at home **2) Follow-up plans for re-evaluation** have been developed and are in place **3) Plans have been developed and are in place in case of change in clinical status**, including means of communication between family and providers and access to emergency medical care.