


**Algorithm for High-Risk Well-Appearing Febrile Infant:
22-60 days of age with bronchiolitis*, temp $\geq 38\text{C}$**



Footnotes:

* AAP **excludes** infants with RSV and / or clinical bronchiolitis or high-risk

**Algorithm for High-Risk Well-Appearing Febrile Infant:
22-60 days of age with bronchiolitis*, temp \geq 38C**

Exclusion criteria

1. Infants who have received immunizations within the last 48 hrs
2. Infants with COVID-19



Consult Pediatrics

Inclusion criteria

- Infants 22-60 days, well-appearing, temp \geq 38C, with clinical bronchiolitis (with or without a positive test for RSV) AND one of the following:
1. Premature infants < 37 wks gestational age
 2. Infants with documented or suspected immune compromise
 3. Infants whose neonatal course was complicated by surgery or infection
 4. Infants with congenital or chromosomal abnormalities
 5. Medically fragile infants requiring some form of technology or ongoing therapeutic intervention to sustain life

CA FIRST Protocol v2.2

Supplemental Information

Footnotes

* AAP **excludes** infants with RSV and / or clinical bronchiolitis or high-risk

Abbreviations

1. Cath: catheter
2. CBC: complete blood count
3. LP: lumbar puncture
4. RSV: respiratory syncytial virus
5. UA: urinalysis
6. UCx: urine culture
7. UTI: urinary tract infection
8. Pabx: parenteral antibiotics